arXiv:1606.03801v1 [math.RA] 13 Jun 2016

Representation Crossed Category of Group-cograded Multiplier Hopf Algebras

Tao Yang *

Abstract Let $A = \bigoplus_{p\in G} A_p$ be a multiplier Hopf T-coalgebra over a group G, in this paper we give the definition of the crossed left A-G-modules and show that the category of crossed left A-G-modules is a monoidal category. Finally we show that a family of multipliers $R = \{R_{p,q} \in M(A_p \otimes A_q)\}_{p,q\in G}$ is a quasitriangular structure of a multiplier T-coalgebra A if and only if the crossed left A-G-module category over A is a braided monoidal category with the braiding c defined by R, generalizing the main results in [14] to the more general framework of multiplier Hopf algebras.
Key words Multiplier Hopf T-coalgebra, quasitriangular, braiding

Mathematics Subject Classification 16W30 · 17B37

1 Introduction

The motivating example for quasitriangular Hopf algebras is given by the Hopf algebra $U_q(\mathrm{g})$ for g a finite-dimensional semisimple Lie algebra over the field $\mathbb{C}$ of complex numbers (see [5]). However, H is not quasitriangular in the strict sense of the definition. The R-matrix lies in a completion of $H \otimes H$ rather than in $H \otimes H$ itself. The $U_q(\mathrm{g})$ is called 'topologically' quasitriangular.

The explicit construction of the universal R-matrix is complicated. There are some ways to evade this problem. One can study the finite-dimensional modules to obtain solutions of the YangBaxter equation. The present approach with multiplier Hopf algebras gives an alternative way to construct a generalized R-matrix in purely algebraic terms. The notion of a quasitriangular (resp. group-cograded) multiplier Hopf algebra is introduced in [13] (resp. [3]).

In 2011, Zhu, Chen and Li [14] studied the relation between the quasitriangular structure of a Hopf group-coalgebra and the braided monoidal category. And in [6], the author considered this relation in the weak Hopf group-coalgebra case. It is natural to ask whether we study the relation between the notion of quasitriangular group-cograded multiplier Hopf algebras and its representation crossed categories.

*Nanjing Agricultural University, Nanjing 210095, Jiangsu, CHINA. E-mail: tao.yang@njau.edu.cn

This question motivates the present research, and in this paper we will give a positive answer to the above question.

The paper is organized in the following way. In section 2, we recall some notions which we will use in the following, such as the definition of multiplier Hopf algebra, multiplier Hopf T-coalgebras and their quasitriangular structures. In section 3, we introduce the notions of (crossed) left A-G-modules, and show that the categories of (crossed) left A-G-modules are both monoidal categories. In section 4, we show the relation between multiplier Hopf T-coalgebra's quasitriangular structure and its representation crossed categories, and prove that the monoidal category $({}_A\mathcal{M}_{crossed}, \otimes, K, a, l, r)$ is a braided monoidal category with the braiding c if and only if A is a quasitriangular multiplier Hopf T-coalgebra, where c is defined by R.

2 Preliminaries

Throughout this paper, all spaces we considered are over a fixed field k. Let A be an (associative) algebra. We do not assume that A has a unit, but we do require that the product, seen as a bilinear form, is non-degenerated. This means that, whenever $a \in A$ and $ab = 0$ for all $b \in A$ or $ba = 0$ for all $b \in A$, we must have that $a = 0$. Then we can consider the multiplier algebra $M(A)$ of A. Recall that $M(A)$ is characterized as the largest algebra with identity containing A as an essential two-sided ideal. In particularly, we still have that, whenever $a \in M(A)$ and $ab = 0$ for all $b \in A$ or $ba = 0$ for all $b \in A$, again $a = 0$. Furthermore, we consider the tensor algebra $A \otimes A$. It is still non-degenerated and we have its multiplier algebra $M(A \otimes A)$. There are natural imbeddings

$$A \otimes A \subseteq M(A) \otimes M(A) \subseteq M(A \otimes A).$$

In generally, when A has no identity, these two inclusions are stict. If A already has an identity, the product is obviously non-degenerate and $M(A) = A$ and $M(A \otimes A) = A \otimes A$. More details about the concept of the multiplier algebra of an algebra, we refer to [8].

Let A and B be non-degenerate algebras, if homomorphism $f : A \longrightarrow M(B)$ is non-degenerated (i.e., $f(A)B = B$ and $Bf(A) = B$), then has a unique extension to a homomorphism $M(A) \longrightarrow M(B)$, we also denote it f.

2.1 Multiplier Hopf algebras

Now, we recall the definition of a multiplier Hopf algebra (see [8] for details). A comultiplication on algebra A is a homomorphism $\Delta : A \longrightarrow M(A \otimes A)$ such that $\Delta(a)(1 \otimes b)$ and $(a \otimes 1)\Delta(b)$ belong to $A \otimes A$ for all $a, b \in A$. We require Δ to be coassociative in the sense that

$$(a \otimes 1 \otimes 1)(\Delta \otimes \iota)(\Delta(b)(1 \otimes c)) = (\iota \otimes \Delta)((a \otimes 1)\Delta(b))(1 \otimes 1 \otimes c)$$

for all $a, b, c \in A$ (where ι denotes the identity map).

A pair (A, Δ) of an algebra A with non-degenerate product and a comultiplication Δ on A is called a *multiplier Hopf algebra*, if the linear map T_1, T_2 defined by

$$T_1(a \otimes b) = \Delta(a)(1 \otimes b), \qquad T_2(a \otimes b) = (a \otimes 1)\Delta(b) \tag{2.1}$$

are bijective.

The bijectivity of the above two maps is equivalent to the existence of a counit and an antipode S satisfying (and defined by)

$$(\varepsilon \otimes \iota)(\Delta(a)(1 \otimes b)) = ab, \qquad m(S \otimes \iota)(\Delta(a)(1 \otimes b)) = \varepsilon(a)b, \tag{2.2}$$

$$(\iota \otimes \varepsilon)((a \otimes 1)\Delta(b)) = ab, \qquad m(\iota \otimes S)((a \otimes 1)\Delta(b)) = \varepsilon(b)a, \tag{2.3}$$

where $\varepsilon : A \longrightarrow k$ is a homomorphism, $S : A \longrightarrow M(A)$ is an anti-homomorphism and m is the multiplication map, considered as a linear map from $A \otimes A$ to A and extended to $M(A) \otimes A$ and $A \otimes M(A)$.

A multiplier Hopf algebra (A, Δ) is called *regular* if (A, Δ^{cop}) is also a multiplier Hopf algebra, where Δ^{cop} denotes the co-opposite comultiplication defined as $\Delta^{cop} = \tau \circ \Delta$ with τ the usual flip map from $A \otimes A$ to itself (and extended to $M(A \otimes A)$). In this case, $\Delta(a)(b \otimes 1), (1 \otimes a)\Delta(b) \in A \otimes A$ for all $a, b \in A$. By Proposition 2.9 in [9], multiplier Hopf algebra (A, Δ) is regular if and only if the antipode S is bijective from A to A.

Remark that any Hopf algebra is a multiplier Hopf algebra and conversely, any multiplier Hopf algebra, with identity, is a Hopf algebra. We will use the adapted Sweedler notation for multiplier Hopf algebras (see [10]), e.g., write $a_{(1)} \otimes a_{(2)}b$ for $\Delta(a)(1 \otimes b)$ and $ab_{(1)} \otimes b_{(2)}$ for $(a \otimes 1)\Delta(b)$.

Let A be a regular multiplier Hopf algebra. Suppose X is a left A-module with the module structure map $\cdot : A \otimes X \longrightarrow X$. We will always assume that the module is non-degenerate, this means that $x = 0$ if $x \in X$ and $a \cdot x = 0$ for all $a \in A$. If the module is unital (i.e., $A \cdot X = X$), then we can get an extension of the module structure to $M(A)$, this means that we can define $f \cdot x$, where $f \in M(A)$ and $x \in X$. In fact, since $x \in X = A \cdot X$, then $x = \sum_i a_i \cdot x_i$, $f \cdot v = \sum_i (fa_i) \cdot x_i$. In this setting, we can easily get $1_{M(A)} \cdot x = x$.

2.2 Multiplier Hopf T-coalgebras and their quasitriangular structures

Let (A, Δ) be a multiplier Hopf algebra and G a group with unit e. Assume that there is a family of (non-trivial) subalgebras $\{A_p\}_{p \in G}$ of A so that

(1) $A = \bigoplus_{p \in G} A_p$ with $A_p A_q = 0$ whenever $p, q \in G$ and $p \neq q$.

(2) $\Delta(A_{pq})(1 \otimes A_q) = A_p \otimes A_q$ and $(A_p \otimes 1)\Delta(A_{pq}) = A_p \otimes A_q$ for all $p, q \in G$.

Then (A, Δ) is called a *G-cograded multiplier Hopf algebra*(see [3, 11]).

We extend the Sweedler notation for a comultiplication in the following way: for any $p, q \in G$, $a \in A_{pq}$ and $a' \in A_q$, we write

$$\Delta_{p,q}(a)(1 \otimes a') = a_{(1,p)} \otimes a_{(2,q)}a'.$$

Let A be a G-cograded multiplier Hopf algebra, then A has the form $A = \bigoplus_{p\in G} A_p$. Assume that there is a group homomorphism $\pi : G \longrightarrow Aut(A)$. We call π an *admissible* action of G on A if also the following requirements hold

(1) $\Delta(\pi_p(a)) = (\pi_p \otimes \pi_p)\Delta(a)$ for all $a \in A$.

(2) $\pi_p(A_q) = A_{\rho_p(q)}$, where ρ is an action of the group G on itself.

(3) $\pi_{\rho_p(q)} = \pi_{pqp^{-1}}$.
This means that the map π takes care of ρ not being the adjoint action. If ρ is the adjoint action, π is called a *crossing*.

A group-cograded multiplier Hopf algebra $A = \bigoplus_{p\in G} A_p$ is said to be a *multiplier Hopf T-coalgebra* provided it is endowed with a crossing π such that each π_q preserves the comultiplication and the counit, i.e., for all $p, q, r \in G$,

$$(\pi_q \otimes \pi_q)\Delta_{p,r} = \Delta_{qpq^{-1},qrq^{-1}}\pi_q, \quad \varepsilon\pi_q = \varepsilon,$$

and π is multiplicative in the sense that $\pi_{pq} = \pi_p\pi_q$ for all $p, q \in G$. It can be considered as generalization of crossed Hopf G-coalgebra introduced in [7].

Let A be a multiplier Hopf T-coalgebra, then we can construct a new regular multiplier Hopf algebra on A by deforming the comultiplication while the algebra structure on A is kept (see Theorem 3.11 in [2]). The comultiplication deformation of A depends on the crossing π in the following way: for all $a \in A$ and $a' \in A_q$,

$$\widetilde{\Delta}(a)(1 \otimes a') = (\pi_{q^{-1}} \otimes \iota)(\Delta(a)(1 \otimes a')).$$

Recall from [3], a G-cograded multiplier Hopf algebra with a crossing action π is called *quasitriangular* if there is a multiplier $R = \sum_{p,q\in G} R_{p,q}$ with $R_{p,q} \in M(A_p \otimes A_q)$ such that

$$\begin{aligned} (\pi_p \otimes \pi_p)(R) &= R, \qquad R\Delta(a) = (\widetilde{\Delta})^{cop}(a)R, \\ (\widetilde{\Delta} \otimes \iota)(R) &= R_{13}R_{23}, \qquad (\iota \otimes \Delta)(R) = R_{13}R_{12}. \end{aligned}$$

for all $p \in G$ and $a \in A$. Sometimes we call R a generalized R-matrix.

3 Category of Crossed Left Modules

The concept of a group-cograded multiplier Hopf algebra was introduced by Abd El-hafez et al in [1] as a generalization of Hopf group-coalgebras introduced in [7]. In the following, similar to the Hopf group-coalgebra case, we will introduce the crossed module category for

a G-cograded multiplier Hopf algebra $A = \bigoplus_{p\in G} A_p$.

Definition 3.1 Let A be a G-cograded multiplier Hopf algebra, a left A-G-module is a family $M = \{M_p\}_{p\in G}$ of vector spaces such that M_p is a left unital A_p-module for any $p \in G$. We denote the structure maps of left A_p-module M_p and A-G-module M by $\Gamma_{M_p} : A_p \otimes M_p \to M_p$ and $\Gamma_M = \{\Gamma_{M_p}\}_{p\in G}$, respectively. Furthermore, if $\{M_p\}_{p\in G}$ and $\{N_p\}_{p\in G}$ are two left A-G-modules, a left A-G-morphism is a family $f = \{f_p : M_p \to N_p\}_{p\in G}$ of linear maps such that f_p is an left A_p-morphism for any $p \in G$.

Denote by ${}_A\mathcal{M}$ the category of all left A-G-modules and its morphisms are left A-G-morphisms. Let $\{M_p\}_{p\in G}$ and $\{N_p\}_{p\in G}$ be two left A-G-modules, we can easily check that $M_s \otimes N_t$ is a unital $A_s \otimes A_t$-module, and a unital left A_{st}-module with the action

$$\Gamma_{M_s\otimes N_t}(a \otimes (m \otimes n)) := a \cdot (m \otimes n) = a_{(1,s)} \cdot m \otimes a_{(2,t)} \cdot n,$$

for all $a \in A_{st}$, $m \in M_s$ and $n \in N_t$. Hence, $(M \otimes N)_p := \bigoplus_{st=p} M_s \otimes N_t$ is a unital left A_p-module. Thus $M \otimes N = \{(M \otimes N)_p\}_{p\in G}$ is a left A-G-module with the structure map $\Gamma_{M\otimes N} = \{\Gamma_{(M\otimes N)_p}\}_{p\in G}$ given by

$$\Gamma_{(M\otimes N)_p} = \bigoplus_{st=p} (\Gamma_{M_s} \otimes \Gamma_{N_t})(\iota \otimes \tau \otimes \iota_{N_t})(\Delta_{s,t} \otimes \iota_{M_s} \otimes \iota_{N_t}). \tag{3.1}$$

Suppose that $L = \{L_p\}_{p\in G}$ is also a left A-G-module, then we have two left A-G-modules $(L \otimes M) \otimes N$ and $L \otimes (M \otimes N)$. By the A-G-module action (3.1), we have for any $p \in G$

$$\begin{aligned}
((L \otimes M) \otimes N)_p &= \bigoplus_{st=p} (L \otimes M)_s \otimes N_t = \bigoplus_{st=p} (\bigoplus_{qr=s} L_q \otimes M_r) \otimes N_t \\
&= \bigoplus_{qrt=p} (L_q \otimes M_r) \otimes N_t, \\
(L \otimes (M \otimes N))_p &= \bigoplus_{qs=p} L_q \otimes (M \otimes N)_s = \bigoplus_{qs=p} L_q \otimes (\bigoplus_{rt=s} M_r \otimes N_t) \\
&= \bigoplus_{qrt=p} L_q \otimes (M_r \otimes N_t).
\end{aligned}$$

One can easily check that $a_{q,r,t} : (L_q \otimes M_r) \otimes N_t \to L_q \otimes (M_r \otimes N_t), (l \otimes m) \otimes n \mapsto l \otimes (m \otimes n)$ is an isomorphism of unital A_{qrt}-modules. Then for any $p \in G$, $a_p = \bigoplus_{qrt=p} a_{q,r,t}$ is an isomorphism of left A_p-modules from $((L \otimes M) \otimes N)_p$ to $(L \otimes (M \otimes N))_p$, and $a = \{a_p\}_{p\in G} : (L \otimes M) \otimes N \to L \otimes (M \otimes N)$ is a left A-G-module isomorphism, it is a family of natural isomorphisms.

Since A_e is a multiplier Hopf algebra, the basic field k is a unital left A_e-module with the action $a \cdot \kappa = \varepsilon(a)\kappa$, where $a \in A_e$ and $\kappa \in k$. Let $K = \{K_p\}_{p\in G}$ with $K_e = k$ and $K_p = 0$ whenever $p \neq e$, then K is a left A-G-module. For any left A-G-module M, $(K \otimes M)_p = K_e \otimes M_p = k \otimes M_p$ and $(M \otimes K)_p = M_p \otimes K_e = M_p \otimes k$. So we have natural

isomorphisms $l_M : K \otimes M \to M$ and $r_M : M \otimes K \to M$ defined by

$$(l_M)_p : k \otimes M_p \to M_p, \qquad \kappa \otimes m \mapsto m,$$
$$(r_M)_p : M_p \otimes k \to M_p, \qquad m \otimes \kappa \mapsto m.$$

That is, $\{l_M\}$ and $\{r_M\}$ are two families of natural isomorphisms of left A-G-modules.

From the above discussion, We summarize the following result:

Theorem 3.2 $({}_A\mathcal{M}, \otimes, K, a, l, r)$ is a monoidal category with the unit object K.

In the following, we will consider the crossed module of multiplier Hopf T-coalgebra $A = \bigoplus_{p\in G} A_p$. Firstly, we introduce the definition of crossed left A-G-module, which generalizes the notion in [14].

Definition 3.3 Let A be a multiplier Hopf T-coalgebra, a left A-G-module $M = \{M_p\}_{p\in G}$ is called crossed if there exists a family $\pi_M = \{\pi_{M,p} : M_q \to M_{pqp^{-1}}\}_{p\in G}$ of linear maps such that the following conditions are satisfied

(1) each $\pi_{M,p} : M_q \to M_{pqp^{-1}}$ is a vector space isomorphism,

(2) each $\pi_{M,p}$ preserves the action, ie., for any $p, q \in G$, $\pi_{M,p}(a \cdot m) = \pi_p(a) \cdot \pi_{M,p}(m)$,

(3) π_M is multiplicative in the sense that for any $p, q \in G$, $\pi_{M,p} \circ \pi_{M,q} = \pi_{M,pq}$.

And for two left crossed A-G-modules $M = \{M_p\}_{p\in G}$ and $N = \{N_p\}_{p\in G}$, a left A-G-module morphism $f = \{f_p\}_{p\in G} : M \to N$ is called crossed if furthermore for any $p, q \in G$, $\pi_{N,q} f_p = f_{qpq^{-1}} \pi_{M,q}$.

Let $A = \bigoplus_{p\in G} A_p$ be a multiplier Hopf T-coalgebra. Denote by ${}_A\mathcal{M}_{crossed}$ the category of all crossed left A-G-modules and crossed left A-G-morphisms. Let M and N be two crossed left A-G-modules, we have already shown that $M \otimes N$ is also a left A-G-module. Define $\pi_{M\otimes N,q} : (M \otimes N)_p \to (M \otimes N)_{qpq^{-1}}$ by

$$\pi_{M\otimes N,q}|_{M_s\otimes N_t} = \pi_{M,q}|_{M_s} \otimes \pi_{N,q}|_{N_t}, \tag{3.2}$$

where $p, q, s, t \in G$ with $p = st$. Since

$$\begin{aligned}
(M \otimes N)_p &= \bigoplus_{st=p} M_s \otimes N_t, \quad \text{and} \\
(M \otimes N)_{qpq^{-1}} &= \bigoplus_{qstq^{-1}=qpq^{-1}} M_{qsq^{-1}} \otimes N_{qtq^{-1}} \\
&= \bigoplus_{st=p} M_{qsq^{-1}} \otimes N_{qtq^{-1}},
\end{aligned}$$

$\pi_{M\otimes N,q}$ is well-defined k-linear isomorphism from $(M \otimes N)_p$ to $(M \otimes N)_{qpq^{-1}}$. Moreover, for $a \in A_p$, $m \in M_s$ and $n \in N_t$ we have $\pi_{M\otimes N,q}(a \cdot (m \otimes n)) = \pi_q(a) \cdot \pi_{M\otimes N,q}(m \otimes n)$.

Indeed,

$$\begin{aligned}\pi_{M\otimes N,q}(a\cdot(m\otimes n)) &= (\pi_{M,q}\otimes\pi_{N,q})(a_{(1,s)}\cdot m\otimes a_{(2,t)}\cdot n)\\ &= \pi_{M,q}(a_{(1,s)}\cdot m)\otimes\pi_{N,q}(a_{(2,t)}\cdot n)\\ &= \pi_q(a_{(1,s)})\cdot\pi_{M,q}(m)\otimes\pi_q(a_{(2,t)})\cdot\pi_{N,q}(n),\end{aligned}$$

and

$$\begin{aligned}\pi_q(a)\cdot\pi_{M\otimes N,q}(m\otimes n) &= \pi_q(a)\cdot(\pi_{M,q}(m)\otimes\pi_{N,q}(n))\\ &= \Delta_{qsq^{-1},qtq^{-1}}(\pi_q(a))\cdot(\pi_{M,q}(m)\otimes\pi_{N,q}(n))\\ &= (\pi_q\otimes\pi_q)\Delta_{s,t}(a)\cdot(\pi_{M,q}(m)\otimes\pi_{N,q}(n))\\ &= \pi_q(a_{(1,s)})\cdot\pi_{M,q}(m)\otimes\pi_q(a_{(2,t)})\cdot\pi_{N,q}(n).\end{aligned}$$

It is easy to get that $\pi_{M\otimes N,q}\circ\pi_{M\otimes N,q'}=\pi_{M\otimes N,qq'}$, i.e., $\pi_{M\otimes N,q}$ is multiplicative. Thus $M\otimes N$ is a crossed left A-G-module.

Now let L be another crossed left A-G-module, one can easily check that

$$\pi_{L\otimes(M\otimes N),q}a_p=a_{qpq^{-1}}\pi_{(L\otimes M)\otimes N,q},$$

and so $a:(L\otimes M)\otimes N\to L\otimes(M\otimes N)$ is a crossed left A-G-module morphism.

Note that $K_e=k=K_{qeq^{-1}}$ and $K_p=0=K_{qpq^{-1}}$ for $p\neq e$. Since $\pi_e:A_p\to A_p$ is the identity map, set $\pi_{K,q}=\iota:K_p\to K_{qpq^{-1}}$, then the unit object $K=\{K_p\}_{p\in G}$ is a crossed left A-G-module.

Finally, one can easily check that the left and right unit constraints $l=\{l_M\}$ and $r=\{r_M\}$ are crossed left A-G-module morphisms.

Therefore we get another main result of this section:

Theorem 3.4 $({}_A\mathcal{M}_{crossed},\otimes,K,a,l,r)$ is a monoidal category with the unit object K.

4 Braided Monnidal Category

Throughout the following, Assume that $A=\bigoplus_{p\in G}A_p$ be a multiplier Hopf T-coalgebra and $R=\{R_{p,q}\in M(A_p\otimes A_q)\}_{p,q\in G}$ be a family of multipliers. Let M and N be any two crossed left A-G-modules, for $s,t\in G$ we define

$$c_{M_s,N_t}:M_s\otimes N_t\longrightarrow N_{sts^{-1}}\otimes M_s$$

by

$$c_{M_s,N_t}(m\otimes n)=(\pi_{N,s}\otimes\iota_{M_s})\tau_{s,t}\Big(R_{s,t}\cdot(m\otimes n)\Big),\tag{4.1}$$

where $m\in M_s$ and $n\in N_t$. For any $p\in G$, define

$$(c_{M,N})_p:(M\otimes N)_p=\bigoplus_{st=p}M_s\otimes N_t\longrightarrow(N\otimes M)_p=\bigoplus_{st=p}N_{sts^{-1}}\otimes M_s$$

by

$$(c_{M,N})_p = \bigoplus_{st=p} c_{M_s,N_t}.$$

Then it is obvious that $(c_{M,N})_p$ is a k-linear isomorphism for any $p \in G$ if and only if so is c_{M_s,N_t} for any $s,t \in G$ with $st = p$.

Lemma 4.1 With the notations above, we have

(1) For any $p \in G$, $(c_{M,N})_p$ is a k-linear isomorphism for any crossed left A-G-modules M and N if and only if R is a family of invertible multipliers.

(2) For any crossed left A-G-modules M and N, $c_{M,N} : M \otimes N \to N \otimes M$ is a left A-G-module morphism if and only if

$$R_{s,t}\Delta_{s,t}(a) = \Big(\tau_{t,s}(\pi_{s^{-1}} \otimes \iota)\Delta_{sts^{-1},s}(a)\Big)R_{s,t} \tag{4.2}$$

for $s,t \in G$ and $a \in A_{st}$.

Proof (1) If $R = \{R_{p,q} \in M(A_p \otimes A_q)\}_{p,q\in G}$ is a family of invertible multipliers, then obviously c_{M_s,N_t} is a k-linear isomorphism for any $s,t \in G$ since so is $\pi_{N,s}$. Conversely, let $M = N = A$, because π_p is an isomorphism, from the hypothesis one knows that the map $A_s \otimes A_t \to A_s \otimes A_t, x \otimes y \mapsto R_{s,t}(x \otimes y)$ is a k-linear isomorphism for any $s,t \in G$. It follows that R is a invertible multiplier in the multiplier algebra $M(A_s \otimes A_t)$.

(2) It is clear that $(c_{M,N})_p$ is an A_p-module morphism if and only if c_{M_s,N_t} is an A_{st}-module morphism for any $s,t \in G$ with $st = p$. Let $m \in M_s, n \in N_t$ and $a \in A_{st}$, then we have

$$\begin{aligned}
c_{M_s,N_t}(a \cdot (m \otimes n)) &= c_{M_s,N_t}(\Delta_{s,t}(a) \cdot (m \otimes n)) \\
&\overset{(4.1)}{=} (\pi_{N,s} \otimes \iota_{M_s})\tau_{s,t}\Big(R_{s,t} \cdot (\Delta_{s,t}(a) \cdot (m \otimes n))\Big) \\
&= (\pi_{N,s} \otimes \iota_{M_s})\tau_{s,t}\Big(R_{s,t}\Delta_{s,t}(a) \cdot (m \otimes n)\Big),
\end{aligned}$$

and

$$\begin{aligned}
a \cdot c_{M_s,N_t}(m \otimes n) &= \Delta_{sts^{-1},s}(a) \cdot c_{M_s,N_t}(m \otimes n) \\
&\overset{(4.1)}{=} (\pi_s \otimes \iota)\Big((\pi_{s^{-1}} \otimes \iota)\Delta_{sts^{-1},s}(a)\Big) \cdot (\pi_{N,s} \otimes \iota_{M_s})\tau_{s,t}\Big(R_{s,t} \cdot (m \otimes n)\Big) \\
&= (\pi_{N,s} \otimes \iota_{M_s})\bigg((\pi_{s^{-1}} \otimes \iota)\Delta_{sts^{-1},s}(a) \cdot \tau_{s,t}\Big(R_{s,t} \cdot (m \otimes n)\Big)\bigg) \\
&= (\pi_{N,s} \otimes \iota_{M_s})\tau_{s,t}\bigg(\tau_{t,s}(\pi_{s^{-1}} \otimes \iota)\Delta_{sts^{-1},s}(a) \cdot \Big(R_{s,t} \cdot (m \otimes n)\Big)\bigg) \\
&= (\pi_{N,s} \otimes \iota_{M_s})\tau_{s,t}\bigg(\Big(\tau_{t,s}(\pi_{s^{-1}} \otimes \iota)\Delta_{sts^{-1},s}(a)\Big)R_{s,t} \cdot (m \otimes n)\bigg).
\end{aligned}$$

Because $\pi_{N,s}$ is an isomorphism, $c_{M_s,N_t}(a\cdot(m\otimes n))=a\cdot c_{M_s,N_t}(m\otimes n)$ if and only if $R_{s,t}\Delta_{s,t}(a)\cdot(m\otimes n)=\Big(\tau_{t,s}(\pi_{s^{-1}}\otimes\iota)\Delta_{sts^{-1},s}(a)\Big)R_{s,t}\cdot(m\otimes n)$. Hence if $R_{s,t}\Delta_{s,t}(a)=\Big(\tau(\pi_{s^{-1}}\otimes\iota)\Delta_{sts^{-1},s}(a)\Big)R_{s,t}$, then c_{M_s,N_t} is an A_{st}-module morphism. Conversely, let $M=N=A$, then for any $x\in A_s$ and $y\in A_t$, $R_{s,t}\Delta_{s,t}(a)(x\otimes y)=\Big(\tau_{t,s}(\pi_{s^{-1}}\otimes\iota)\Delta_{sts^{-1},s}(a)\Big)R_{s,t}(x\otimes y)$. By the non-degenerate of the product, we can get $R_{s,t}\Delta_{s,t}(a)=\Big(\tau_{t,s}(\pi_{s^{-1}}\otimes\iota)\Delta_{sts^{-1},s}(a)\Big)R_{s,t}$. □

Lemma 4.2 The following two statements are equivalent:

(1) $\pi_{N\otimes M,q}\circ(c_{M,N})_p=(c_{M,N})_{qpq^{-1}}\circ\pi_{M\otimes N,q}$ for any crossed left A-G-modules M and N, and $p,q\in G$.

(2) $(\pi_q\otimes\pi_q)R_{s,t}=R_{qsq^{-1},qtq^{-1}}$ for any $q,s,t\in G$.

Proof For any $s,t,q\in G$, $m\in M_s$ and $n\in N_t$,

$$\begin{aligned}
&\pi_{N\otimes M,q}\circ(c_{M,N})_{st}(m\otimes n)\\
=\;&(\pi_{N,q}\otimes\pi_{M,q})(c_{M_s,N_t})(m\otimes n)\\
\overset{(4.1)}{=}\;&(\pi_{N,q}\otimes\pi_{M,q})(\pi_{N,s}\otimes\iota_{M_s})\tau_{s,t}\Big(R_{s,t}\cdot(m\otimes n)\Big)\\
=\;&(\pi_{N,qs}\otimes\pi_{M,q})\Big(\tau_{s,t}(R_{s,t})\cdot(n\otimes m)\Big)\\
=\;&(\pi_{N,qsq^{-1}}\otimes\iota_{M_{qsq^{-1}}})(\pi_{N,q}\otimes\pi_{M,q})\Big(\tau_{s,t}(R_{s,t})\cdot(n\otimes m)\Big)\\
=\;&(\pi_{N,qsq^{-1}}\otimes\iota_{M_{qsq^{-1}}})\Big((\pi_q\otimes\pi_q)(\tau_{s,t}R_{s,t})\cdot(\pi_{N,q}(n)\otimes\pi_{M,q}(m))\Big)\\
=\;&(\pi_{N,qsq^{-1}}\otimes\iota_{M_{qsq^{-1}}})\left(\tau_{qsq^{-1},qtq^{-1}}(\pi_q\otimes\pi_q)(R_{s,t})\cdot\Big(\pi_{N,q}(n)\otimes\pi_{M,q}(m)\Big)\right),
\end{aligned}$$

and

$$\begin{aligned}
&(c_{M,N})_{qstq^{-1}}\circ\pi_{M\otimes N,q}(m\otimes n)\\
=\;&c_{M_{qsq^{-1}},N_{qtq^{-1}}}\big(\pi_{M,q}(m)\otimes\pi_{N,q}(n)\big)\\
\overset{(4.1)}{=}\;&(\pi_{N,qsq^{-1}}\otimes\iota_{M_{qsq^{-1}}})\tau_{qsq^{-1},qtq^{-1}}\Big(R_{qsq^{-1},qtq^{-1}}\cdot\big(\pi_{M,q}(m)\otimes\pi_{N,q}(n)\big)\Big)\\
=\;&(\pi_{N,qsq^{-1}}\otimes\iota_{M_{qsq^{-1}}})\Big(\tau_{qsq^{-1},qtq^{-1}}(R_{qsq^{-1},qtq^{-1}})\cdot\big(\pi_{N,q}(n)\otimes\pi_{M,q}(m)\big)\Big).
\end{aligned}$$

It follows that (2) implies (1). Conversely, assume that (1) is satisfied. Because $\pi_{N,qsq^{-1}}$ is an isomorphism, it follows $(\pi_q\otimes\pi_q)(R_{s,t})\cdot(\pi_{M,q}(m)\otimes\pi_{N,q}(n))=R_{qsq^{-1},qtq^{-1}}\cdot(\pi_{M,q}(m)\otimes\pi_{N,q}(n))$. Because $\pi_{M,q}$ and $\pi_{N,q}$ are isomorphisms, $(\pi_q\otimes\pi_q)(R_{s,t})\cdot(u\otimes v)=R_{qsq^{-1},qtq^{-1}}\cdot(u\otimes v)$ for any $u\in M_{qsq^{-1}}$ and $v\in N_{qtq^{-1}}$. Let $M=N=A$, then by the non-degenerate of the product, we can get $(\pi_q\otimes\pi_q)(R_{s,t})=R_{qsq^{-1},qtq^{-1}}$. □

Lemma 4.3 The following two statements hold:

(1) $c_{L,M\otimes N}=(\iota_M\otimes c_{L,N})(c_{L,M}\otimes\iota_N)$ for any crossed left A-G-modules L, M and N, if and only if for any $r,s,t\in G$,

$$(\iota\otimes\Delta_{s,t})(R_{r,st})=(R_{r,t})_{1s3}(R_{r,s})_{12t}.$$

(2) $c_{L\otimes M,N}=(c_{L,N}\otimes\iota_M)(\iota_L\otimes c_{M,N})$ for any crossed left A-G-modules L, M and N, if and only if for any $r,s,t\in G$,

$$(\Delta_{r,s}\otimes\iota)(R_{rs,t})=\Big((\iota\otimes\pi_{s^{-1}})R_{r,sts^{-1}}\Big)_{1s3}(R_{s,t})_{r23}.$$

Proof We only check the first statement and the second one is similar. For $v\in L_r$, $m\in M_s$ and $n\in N_t$,

$$\begin{aligned}
&(c_{L,M\otimes N})_{rst}(v\otimes m\otimes n)\\
=\;&(\pi_{M\otimes N,r}\otimes\iota_{L_r})\tau_{r,st}\Big(R_{r,st}\cdot(v\otimes(m\otimes n))\Big)\\
=\;&(\pi_{M\otimes N,r}\otimes\iota_{L_r})(\iota_{M_s}\otimes\tau_{r,t})(\tau_{r,s}\otimes\iota_{N_t})\Big((\iota\otimes\Delta_{s,t})R_{r,st}\cdot(v\otimes m\otimes n)\Big),
\end{aligned}$$

and

$$\begin{aligned}
&(\iota_M\otimes c_{L,N})(c_{L,M}\otimes\iota_N)(v\otimes m\otimes n)\\
\overset{(4.1)}{=}\;&(\iota_M\otimes c_{L,N})\Big((\pi_{M,r}\otimes\iota_{L_r})\tau_{r,s}(R_{r,s}\cdot(v\otimes m))\otimes n\Big)\\
=\;&(\pi_{M,r}\otimes c_{L,N})(\tau_{r,s}\otimes\iota_{N_t})\Big((R_{r,s})_{12t}\cdot(v\otimes m\otimes n)\Big)\\
=\;&(\pi_{M,r}\otimes c_{L,N})(R^2_{r,s}\cdot m\otimes R^1_{r,s}\cdot v\otimes n)\\
\overset{(4.1)}{=}\;&(\pi_{M,r}\otimes(\pi_{N,r}\otimes\iota_{L_r}))(\iota_{M_s}\otimes\tau_{r,t})\Big(R^2_{r,s}\cdot m\otimes R^1_{r,t}R^1_{r,s}\cdot v\otimes R^2_{r,t}\cdot n\Big)\\
=\;&(\pi_{M,r}\otimes(\pi_{N,r}\otimes\iota_{L_r}))(\iota_{M_s}\otimes\tau_{r,t})(\tau_{r,s}\otimes\iota_{N_t})\Big(R^1_{r,t}R^1_{r,s}\cdot v\otimes R^2_{r,s}\cdot m\otimes R^2_{r,t}\cdot n\Big)\\
=\;&(\pi_{M,r}\otimes(\pi_{N,r}\otimes\iota_{L_r}))(\iota_{M_s}\otimes\tau_{r,t})(\tau_{r,s}\otimes\iota_{N_t})\Big((R_{r,t})_{1s3}(R_{r,s})_{12t}\cdot(v\otimes m\otimes n)\Big)\\
\overset{(4.1)}{=}\;&(\pi_{M\otimes N,r}\otimes\iota_{L_r})(\iota_{M_s}\otimes\tau_{r,t})(\tau_{r,s}\otimes\iota_{N_t})\Big((R_{r,t})_{1s3}(R_{r,s})_{12t}\cdot(v\otimes m\otimes n)\Big).
\end{aligned}$$

Thus if $(\iota\otimes\Delta_{s,t})(R_{r,st})=(R_{r,t})_{1s3}(R_{r,s})_{12t}$, then $c_{L,M\otimes N}=(\iota_M\otimes c_{L,N})(c_{L,M}\otimes\iota_N)$. Conversely, since $\pi_{M\otimes N,r}$, $\tau_{r,t}$ and $\tau_{r,s}$ are bijective, $(\iota\otimes\Delta_{s,t})R_{r,st}\cdot(v\otimes m\otimes n)=(R_{r,t})_{1s3}(R_{r,s})_{12t}\cdot(v\otimes m\otimes n)$. Let $L=M=N=A$, at this time the module action is the product of A, because this product is non-degenerate, we can easily get $(\iota\otimes\Delta_{s,t})R_{r,st}=(R_{r,t})_{1s3}(R_{r,s})_{12t}$. □

Theorem 4.4 Lat $A=\bigoplus_{p\in G}A_p$ be a multiplier Hopf T-coalgebra and $R=\{R_{p,q}\in M(A_p\otimes A_q)\}_{p,q\in G}$ be a family of multipliers. Then the monoidal category $({}_A\mathcal{M}_{crossed},\otimes,K,a,l,r)$ is a braided monoidal category with the braiding c if and only if A is a quasitriangular multiplier Hopf T-coalgebra, where c is defined by R as in equation (4.1).

Proof If c is a braiding of the monoidal category, then it follows from Lemmas 4.1, 4.2 and 4.3 that R is a quasitriangular structure. Conversely, assume that R is a quasitriangular structure, then by Lemmas 4.1, 4.2 and 4.3, it is enough to show that $c=\{c_{M,N}\}$ is natural.

Let $f : M \to M'$ and $g : N \to N'$ be two crossed morphisms of crossed left A-G-modules, in the following we need to check the braiding preserves the morphisms, i.e., $(g \otimes f)c_{M,N} = C_{M',N'}(f \otimes g)$. In fact, for any $m \in M_s$ and $n \in N_t$ with $\forall s, t \in G$,

$$\begin{aligned}
((g \otimes f)c_{M,N})_{st}(m \otimes n) &= (g_{sts^{-1}} \otimes f_s)c_{M_s,N_t}(m \otimes n) \\
&= (g_{sts^{-1}} \otimes f_s)(\pi_{N,s} \otimes \iota_{M_s})\tau_{s,t}\big(R_{s,t} \cdot (m \otimes n)\big) \\
&= (g_{sts^{-1}} \otimes f_s)\Big((\pi_{N,s} \otimes \iota_{M_s})\big(\tau_{s,t}R_{s,t} \cdot (n \otimes m)\big)\Big) \\
&= (g_{sts^{-1}} \otimes f_s)\Big((\pi_s \otimes \iota)(\tau_{s,t}R_{s,t}) \cdot (\pi_{N,s} \otimes \iota_{M_s})(n \otimes m)\Big) \\
&= (\pi_s \otimes \iota)(\tau_{s,t}R_{s,t}) \cdot \Big((g_{sts^{-1}} \otimes f_s)(\pi_{N,s} \otimes \iota_{M_s})(n \otimes m)\Big) \\
&= (\pi_s \otimes \iota)(\tau_{s,t}R_{s,t}) \cdot \Big(\pi_{N',s} \circ g_t(n) \otimes f_s(m)\Big),
\end{aligned}$$

and

$$\begin{aligned}
(C_{M',N'}(f \otimes g))_{st}(m \otimes n) &= C_{M'_s,N'_t}(f_s(m) \otimes g_t(n)) \\
&= (\pi_{N',s} \otimes \iota_{M'_s})\tau_{s,t}\Big(R_{s,t} \cdot (f_s(m) \otimes g_t(n))\Big) \\
&= (\pi_{N',s} \otimes \iota_{M'_s})\Big(\tau_{s,t}R_{s,t} \cdot (g_t(n) \otimes f_s(m))\Big) \\
&= (\pi_s \otimes \iota)(\tau_{s,t}R_{s,t}) \cdot \Big((\pi_{N',s} \otimes \iota_{M'_s})(g_t(n) \otimes f_s(m))\Big) \\
&= (\pi_s \otimes \iota)(\tau_{s,t}R_{s,t}) \cdot \Big(\pi_{N',s} \circ g_t(n) \otimes f_s(m)\Big).
\end{aligned}$$

This completes the proof. □

Acknowledgements

The work was partially supported by the NNSF of China (No. 11226070), the NJAUF (No. LXY201201019, LXYQ201201103) and NSF for Colleges and Universities in Jiangsu Province (No. 11KJB110004).

References

[1] Abd El-hafez, A. T., Delvaux, L. and Van Daele, A. (2007). Group-cograded multiplier Hopf (∗-)algebra. *Algebras and Representation Theory* 10: 77-95.

[2] Delvaux, L. and Van Daele, A. (2007). The Drinfel'd Double for group-cograded multiplier Hopf algebras. *Algebras and Representation Theory* 10(3): 197-221.

[3] Delvaux, L., Van Daele, A. and Wang S. H. (2005). Quasitriangular (G-cograded) multiplier Hopf algebras. *Journal of Algebra* 289: 484-514.

[4] Drabant, B., Van Daele, A. and Zhang, Y. (1999). Actions of multiplier Hopf algebras. *Communications in Algebra* 27(9): 4117-4127.

[5] Drinfeld, V.G. (1987). Quantum groups. *Proceedings of the International Congress of Mathematicians* Berkeley, California, USA, 1986: 798-820.

[6] Guo, S. J. (2015). Representation Crossed Category of Weak Hopf Group Coalgebras. *Southeast Asian Bulletin of Mathematics* 39: 487495.

[7] Turaev, V. G. (2008). Crossed group-categories. *Arabian Journal for Science and Engineering* 33(2C): 483-503.

[8] Van Daele, A. (1994). Multiplier Hopf algebras. *Transaction of the American Mathematical Society* 342(2): 917-932.

[9] Van Daele, A. (1998). An algebraic framework for group duality. *Advance in Mathematics* 140(2): 323-366.

[10] Van Daele, A. (2008). Tools for working with multiplier Hopf algebras. *The Arabian Journal for Science and Engineering* 33(2C): 505-527.

[11] Yang, T. and Wang, S. H. (2011). A lot of quasitriangular group-cograded multiplier Hopf algebras. *Algebras and Representation Theory* 14(5): 959-976.

[12] Yang, T.and Wang, S. H. (2011). Constructing new braided T-categories over regular multiplier Hopf algebras. *Communications in Algebra* 39(9): 3073-3089.

[13] Zhang, Y. H. (1999). The quantum double of a coFrobenius Hopf algebra. *Communications in Algebra* 27(3): 1413-1427.

[14] Zhu, M. L., Chen, H. X. and Li, L. B. (2011). Quasitriangular Hopf group coalgebras and braided monoidal categories. *Arabian Journal for Science and Engineering* 36: 10631070.